

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representation of  
The original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

FIGURE 1


FPLC elution profile of the NDS as prepared by the modified method.



09964240-092601

FIGURE 2

Continuous scan (200 - 700 nm) of a typical lot of the NDS as prepared by the modified method.






FIGURE 3






FIGURE 4



The PCE-induced dose-dependent production of cytokines in 48hr cultures of PBMC derived from two different blood donors.

FIGURE 5



Interferon gamma production by allogeneic CD3+ cells upon interaction with increasing numbers of PCE or cytokine treated CD14+ cells. The number of stimulator cells (10000, 2500, 625) produced ratios of stimulators to targets of 10:1, 40:1, 160:1, respectively. Triangles represent cytokine-generated DC, squares represent PCE-generated DC, and diamonds represent media control cells.

FIGURE 6

0964240.092601

## CD8 cell IFN- $\gamma$ production





FIGURE 7.



## CD4 cell IFN- $\gamma$ production



Intracellular interferon gamma production by CD8+ or CD4+ splenocytes isolated from animals vaccinated with irrelevant DNA (pICAT) or with pIGag + oral PCE, pIGag + IM PCE, or pIGag DNA only and then stimulated with the HIV peptide p7g (for the CD8 response) or recombinant p24 (for the CD4 response).

FIGURE 7a